

String field theory Hamiltonians from Yang-Mills theories

Vipul Periwal

Department of Physics, Princeton University, Princeton, New Jersey 08544

Abstract

Marchesini showed that the Fokker–Planck Hamiltonian for Yang–Mills theories is the loop operator. Jevicki and Rodrigues showed that the Fokker–Planck Hamiltonian of some matrix models coincides with temporal gauge non-critical string field theory Hamiltonians constructed by Ishibashi and Kawai (and their collaborators). Thus the loop operator for Yang–Mills theory is the temporal gauge Hamiltonian for a noncritical string field theory, in accord with Polyakov’s conjecture. The consistency condition of the string interpretation is the zigzag symmetry emphasized by Polyakov. Several aspects of the noncritical string theory are considered, relating the string field theory Hamiltonian to the worldsheet description.

Mandelstam [1] realized the importance of gauge invariant loop observables in Yang–Mills theory. See also [2]. The fact that a string interpretation for Yang–Mills theories is natural is particularly transparent in the loop equation derived by Guerra and collaborators [3], and rediscovered in [4]. This Schwinger–Dyson equation governs the dynamics of gauge invariant Wilson loops in Yang–Mills theory. There are simple geometric interpretations for the various terms that appear in terms of string propagation and interactions, with the string joining interaction suppressed by a factor of $1/N^2$ relative to the string splitting interaction in a normalization natural for the large N limit.

The loop equation governing Wilson loop expectation values in gauge theories can be written as the expectation value of the action of an operator, the loop operator, acting on Wilson lines. It is a fundamental observation due to Marchesini [5] that the loop operator coincides with the Fokker–Planck Hamiltonian that appears in the stochastic quantization [6] of the gauge theory. As such, the loop operator plays a central rôle in gauge theories but its significance in, for example, a string field theory equivalent to a Yang–Mills theory has not been elucidated. The contribution of this paper is to point out that *the loop operator is precisely the temporal gauge string field theory Hamiltonian*. I show that the consistency condition for this identification is Polyakov’s zig–zag symmetry. I explain how the peculiar asymmetry between joining and splitting vertices can be accounted for in a worldsheet description, and I explain how recent conjectures on Yang–Mills theories and their dimensional reductions [7–9] are related to the explicit identification given here.

The explicit connection given in this paper between string theories and gauge theories should be compared to the efforts that have gone into attempted constructions of worldsheet descriptions of first–quantized string propagation in supergravity backgrounds [12] believed to be dual to gauge theories. The conceptual simplicity of the string field theory Hamiltonian

is striking because in no other instance in string theory is a string field theory simpler or easier to construct than the first-quantized string theory. The identification of this Hamiltonian can be used to deduce properties of a worldsheet first-quantized description, as I shall show below. It is plausible that the first-quantized description will capture only part of the physics incorporated in the *exact* string field theory Hamiltonian, just as an expansion in Feynman diagrams misses non-perturbative physics.

Concretely, the string field theory that arises from the Yang–Mills theory is a *noncritical* string theory. The spacetime dimension in which the string propagate is therefore one more than the sum of the number of adjoint scalars and the number of gauge potential components. Thus, *e.g.* for the $d = 4$ maximally supersymmetric Yang–Mills theory, the spacetime dimension of the noncritical string theory is 11, which differs from the naïve interpretation of the Maldacena conjecture [8]. However the present construction precisely agrees with the conjectured QCD string in Polyakov’s approach [7] which has been argued on physical grounds to be a noncritical string living in five dimensions. Since the infinite N Maldacena conjecture is supported by several precise comparisons with IIB supergravity in ten dimensions, it is important to explain how a noncritical string theory can maintain this agreement. I explain this below by recalling the difference between noncritical string measures and critical string measures following [10,11]. Furthermore, comparing to the calculation of Dorey, Hollowood, Khoze, Mattis and Vandoren [13], it is interesting to observe that at finite N the instanton calculation does not localize on S^5 but rather seems to involve an 11-dimensional space [14]. (The connection to the noncritical string given in a previous version of this paper was not mentioned in [14].)

The loop operator is easiest understood in the lattice theory since a precise interpretation requires a cutoff—this cutoff is not necessarily the same as the cutoff used in defining the quantum gauge theory [15]. Define the loop function for a $U(N)$ gauge theory on a d -dimensional lattice

$$W(C) \equiv \frac{1}{N} \text{Tr} U(C) \quad (1)$$

where C is a closed contour on the lattice, and $U(C)$ is the path ordered product of link variables along this contour $U_{\mu_1}(x_1)U_{\mu_2}(x_2)\dots U_{\mu_n}(x_n)$. Define an electric field operator appropriate for a $d + 1$ -dimensional theory [5]:

$$E_\mu : \left[E_\mu^a(x), U_\nu(y) \right] = T^a U_\nu(y) \delta_{\mu\nu} \delta_{x,y} \quad (2)$$

where $\mu, \nu = 1, \dots, d$, and $a = 1, \dots, N^2$, with $\text{Tr} T^a T^b = N/2$. For any lattice action, $S(U)$, the stochastic Fokker–Planck Hamiltonian is

$$H \equiv \frac{2}{N} \sum_{x,\mu,a} e^{-S(U)} E_\mu^a(x) e^{S(U)} E_\mu^a(x) ; \quad (3)$$

while H is not Hermitian, by a similarity transformation we can make it Hermitian.

There are simple geometric interpretations for the terms that appear in H , which are clearer when we write

$$H \equiv \frac{2}{N} \sum_{x,\mu,a} E_\mu^a(x) E_\mu^a(x) + \left[E_\mu^a(x), S(U) \right] E_\mu^a(x). \quad (4)$$

The first term, which is independent of the lattice action, includes terms that correspond to the splitting and joining of strings, with the latter term suppressed relative to the former by a factor of N^{-2} . The equilibrium condition for the stochastic field theory correlation function turns out to be the loop equation [5]:

$$\langle HW(C) \rangle = 0. \quad (5)$$

Independently, Jevicki and Sakita [16] showed that the large N saddle point equation of Yang–Mills theory formulated as a collective field theory is equivalent to the loop equation, as well.

The asymmetry in the powers of N for joining and splitting interactions is a clue for the string field theory interpretation since in temporal gauge one does find such an asymmetry between string splitting and string joining vertices [17]. This reassignment of powers of N is consistent with the usual topological loop counting of string diagrams. The S -dependent term has operators corresponding to the motion in the space of loops, and tadpole operators that annihilate loops. Thinking of S as defining the background configuration of the space of loops, the propagation and annihilation of loops is background dependent, but the interactions of loops are background independent. Notice that the 't Hooft coupling $g_{\text{YM}}^2 N$ appears only in this term. This is essentially another realization of an idea of Horowitz, Lykken, Rohm and Strominger [18] regarding string field theory. In the large N limit there are no terms that can join strings. Thus in the large N limit, H acts as a derivation on products $\prod_i W(C_i)$.

In a beautiful set of papers, Ishibashi, Kawai, and collaborators [17,19,20] have constructed non-critical string field theories in temporal gauge. As an example, consider their string field theory Hamiltonian for the $c = 0$ model:

$$H_{\text{IK}} = \int_0^\infty dl_1 dl_2 \Psi_{l_1}^\dagger \Psi_{l_2}^\dagger \Psi_{l_1+l_2} (l_1 + l_2) + \int_0^\infty dl \rho(l) \Psi_l + O(g^2) \quad (6)$$

where Ψ_l annihilates a string of length l , and $[\Psi_l, \Psi_{l'}^\dagger] = \delta(l - l')$. The term of $O(g^2)$ is

$$g^2 \int_0^\infty dl_1 dl_2 \Psi_{l_1+l_2}^\dagger \Psi_{l_2} \Psi_{l_1} l_1 l_2 \quad (7)$$

and describes the merging of strings. Ψ annihilates unmarked loops and $\Psi^\dagger$ creates marked loops. The string joining and string splitting terms have different powers of g , just as there are different powers of N^{-1} in the loop joining and splitting terms in the loop operator. This Hamiltonian was first written down for the $c = 1$ model by Das and Jevicki [21] and it was pointed out subsequently by Moore, Seiberg and Staudacher [22] that it applied as well to $c < 1$ models with changes in the tadpole term.

A connected amplitude with b boundaries and h handles comes with a factor $g^{2h-2+2b}$, which is not the usual topological combination. (I have corrected a small error in [20] here.) This can be traced to the fact that the disk amplitude is non-zero and has a factor g^0 in this formalism. Normalizing connected amplitudes with respect to the conventionally normalized disk amplitudes gives the usual Euler characteristic power $g^{-(2-2h-b)}$. In the temporal gauge interpretation, the tadpole term is $O(1)$, just as for large N Yang–Mills theory the normalized expectation value is $O(1)$. One could rescale Ψ and $\Psi^\dagger$ while preserving

the commutation relation to make the joining and the splitting terms of the same order in g , but this would change the normalization of the tadpole term as well, making the expectation value of a Wilson loop $O(N)$, appropriate for a string disk amplitude. This is not, however, natural from the point of view of the gauge theory, so I will give below another geometric interpretation of the powers of g .

Jevicki and Rodrigues [23] showed that the Hamiltonian H_{IK} (including the $O(g^2)$ term) arises naturally as the double-scaling limit [24] of the $k = 2$ matrix model Fokker–Planck Hamiltonian. Thus one has a direct map from matrix models to noncritical string field theories in the temporal gauge, via the Fokker–Planck Hamiltonian.

Crucial to the string interpretation of the Fokker–Planck Hamiltonian is a consistency check, equivalent to diffeomorphism invariance on the string worldsheet [20,23]. In the case of the loop operator, it is a generalization of the derivation given in [20](Sect. 3) that the consistency of the string interpretation is equivalent to the zigzag symmetry of Wilson loops, particularly emphasized by Polyakov [7]. The key point to note here is that two Wilson loops $L_i, i = 1, 2$ (which are the strings in this formalism) may join either by an infinitesimal loop attaching to L_1 leading to contact with L_2 or vice versa. For the string interpretation, diffeomorphism invariance on the worldsheet implies that the difference between these two amplitudes should vanish. Subtracting the amplitudes for the two processes, we encounter amplitudes involving insertions of infinitesimal back-tracking loops, which are trivial if and only if the zig–zag symmetry holds.

In connection with the zig–zag symmetry, a recent investigation of the loop equation [9] in the AdS/CFT correspondence [25] has found that the agreement between gauge theory expectations and critical string expectations is regularization dependent. Recalling the analysis of Polyakov [7] and Klebanov, Kogan and Polyakov [11] suggests then that the zig–zag symmetry requires the non-critical Liouville dimension to be taken into account.

Thus we reach the main conclusion of this paper: The loop operator of Yang–Mills theory is the temporal gauge string field theory Hamiltonian of a noncritical string theory, provided that the zig–zag symmetry is maintained.

This observation suggests that the appropriate place to look for a worldsheet description of the Yang–Mills string is temporal gauge [19]. This gauge is quite different from conformal gauge so much of the usual intuition for first-quantized string theory requires revision. For example, in light-cone string field theory, for example, string splitting and joining interaction vertices are given by mirror image worldsheets, with curvature singularities precisely at the joining/splitting point. These curvature singularities are integrable and precisely lead to the expected power of g for both string joining and splitting vertices, since the spacetime dilaton Φ couples to $\int {}^{(2)}R$. The incoming and outgoing strings lie on curves with vanishing geodesic curvature in light-cone string field theory. In temporal gauge, as mentioned above, the splitting and joining interactions come with different powers of g . Since the Euler characteristic of the pants diagram is $\chi_E = -1$, how can this be? A simple resolution of this puzzle is to note that only the worldsheet bulk curvature couples to the dilaton, whereas the Euler characteristic is a sum of two terms

$$\int {}^{(2)}R + \sum_{\text{boundaries}} \oint_{C_i} \kappa = 2\pi\chi_E \quad (8)$$

according to Gauss and Bonnet [26], with κ the geodesic curvature of the boundary components. There is a time asymmetry in the definition of temporal gauge, so we can sensibly

distinguish between incoming and outgoing strings. Let $\oint \kappa = 2\pi K_{\pm}$ for ‘standard’ outgoing/incoming loops respectively, then for the joining vertex we have

$$\frac{1}{2\pi} \int {}^{(2)}R = -2 = \chi_E(\text{pants}) - K_+ - 2K_- \quad (9)$$

and for the splitting vertex we have

$$\frac{1}{2\pi} \int {}^{(2)}R = 0 = \chi_E(\text{pants}) - 2K_+ - K_- . \quad (10)$$

We find then

$$K_+ = -K_- = -1. \quad (11)$$

This is *precisely* as it must be since in the standard gluing of worldsheet diagrams in string field theory, the geodesic curvatures of boundaries being identified must cancel for constructing smooth surfaces. As an independent check this predicts that the tadpole diagram (a disk) will have one incoming string giving a contribution $K_- = +1$ to the Euler characteristic and therefore have vanishing integrated bulk curvature,

$$\chi_E(\text{disk}) - K_- = +1 - 1 = 0 = \frac{1}{2\pi} \int {}^{(2)}R \quad (12)$$

implying that the tadpole will be $O(1)$, which is exactly correct. As another consistency check, the Euler characteristic of the cylinder is zero, thus incoming and outgoing boundaries must have boundary geodesic curvature that differs only in sign since the propagation term in the loop operator has no factors of N and therefore corresponds to a vanishing bulk contribution. The fact that there is nonvanishing geodesic curvature associated with the boundaries in this gauge is related to the fact that the Wilson loops are normalized with factors of $1/N$.

The discussion trivially extends to dimensionally reduced and/or supersymmetric [27] gauge theories, with $A_{\mu}(x(s))dx^{\mu}/ds$ replaced by $\Phi_i(x(s))\sum_n v^i\delta(s-s_n)$ in the path-ordered exponential for dimensionally reduced directions. This is of interest in light of the conjectures of Banks, Fischler, Shenker and Susskind [28], and Maldacena [8], though it needs to be emphasized again that these conjectures are associated with critical string theories.

In Polyakov’s attempts at constructing string theories equivalent to Yang–Mills theory, on the other hand, the strings that appear are non-critical strings [7]. As such, the identification which has been proposed in this paper is exactly in line with his work, especially noting the zig-zag symmetry’s rôle in a string interpretation.

The string field theory that I have identified in this paper is gauge-fixed. This is related to the fact that we have worked entirely in terms of gauge-invariant operations on gauge-invariant Wilson loops. Recollect that residual spacetime gauge transformations in critical string theory appear as conformal transformations on the worldsheet in conformal gauge.

Since I have given here a construction of a non-critical string theory from Yang–Mills theory that is different from the standard conjectures [9], I must explain how the numerous agreements supporting the standard conjectures that appear at string tree-level come about. This is easily done: What is the difference between a critical string amplitude and a noncritical string amplitude? The functional measure for a critical string amplitude has

an integration over the conformal mode and a division by the volume of the group of conformal transformations, $\text{vol}(\text{Conf})$. These cancel for correlation functions of on-shell vertex operators, and do not cancel if we consider off-shell amplitudes [10]. Even for off-shell amplitudes, on the sphere, there are no moduli and $\text{vol}(\text{Conf})$ is a amplitude independent constant. On higher genus surfaces, $\text{vol}(\text{Conf})$ depends on the moduli and cannot be factored out. For noncritical strings, there is still an integration over the conformal mode, but no corresponding division by $\text{vol}(\text{Conf})$. Thus, if we are interested in tree amplitudes, we can extract critical string tree amplitudes from a subset of the noncritical string tree amplitudes. (Recall that string tree amplitudes do not suffice to reconstruct string loop amplitudes [29].) This subset corresponds to on-shell string states that have no dependence on the conformal mode. From the Fokker–Planck perspective, it is natural to focus on correlations

$$\lim_{D \uparrow \infty} \langle 0 | \exp(-DH) \prod W(C_i) | 0 \rangle \quad (13)$$

where D is the Fokker–Planck time (the Liouville dimension coördinate), here identified with the worldsheet time coördinate, of Wilson loops C_i that satisfy the equilibrium condition

$$[H, W(C_i)] = 0 \text{ for each } i. \quad (14)$$

It seems clear that correlations of observables satisfying the equilibrium condition will correspond to amplitudes with truncated external leg propagators, natural for on-shell vertex operators in critical string theory. However, the precise relation between these two sets of objects remains to be computed.

Lastly, I mention that supporting evidence for my identification comes from a very recent preprint by Lidsey [30]. In this work it is shown that the critical IIB supergravity backgrounds can be embedded in eleven-dimensional Ricci-flat spaces, including backgrounds with non-trivial Ramond–Ramond fields. This is precisely in accord with the above-mentioned relation between critical strings and non-critical strings at tree level, see especially Klebanov, Kogan and Polyakov [11].

I am grateful to I. Klebanov, G. Lifschytz, J. Polchinski and A. Polyakov for helpful conversations and A. Jevicki for a useful communication. This work was supported in part by NSF grant PHY-9802484.

REFERENCES

- [1] S. Mandelstam, Ann. of Phys. 19 (1962) 25; Phys. Rev. 175 (1968) 1580
- [2] J.-L. Gervais and A. Neveu, Phys. Lett. B80 (1979) 255; Y. Nambu, Phys. Lett. B80 (1979) 372; A. Polyakov, Phys. Lett. B82 (1979) 247
- [3] G. De Angelis, D. De Falco and F. Guerra, Nuovo Cim. Lett. 19 (1977) 55; F. Guerra, R. Marra and G. Immirzi, Nuovo Cim. Lett. 23 (1978) 237
- [4] D. Foerster, Phys. Lett. 87B (1979) 83; T. Eguchi, Phys. Lett. 87B (1979) 91; Yu. Makeenko and A. Migdal, Phys. Lett. 88B (1979) 135
- [5] G. Marchesini, Nucl. Phys. B191 (1981) 214; B239 (1984) 135
- [6] G. Parisi and Y.-S. Wu, Sci. Sin. 24 (1981) 484
- [7] A. Polyakov, Nucl. Phys. Proc. Supp. 68 (1998) 1; Int. J. Mod. Phys. A14 (1999) 645
- [8] J. Maldacena, Adv. Theor. Math. Phys. 2 (1998) 231; S. Gubser, I. Klebanov and A. Polyakov, Phys. Lett. B428 (1998) 105; E. Witten, Adv. Theor. Math. Phys. 2 (1998) 253
- [9] O. Aharony, S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, *Large N field theories, string theory and gravity*, hep-th/9905111
- [10] R.C. Myers and V. Periwai, Phys. Rev. Lett. 70 (1993) 2841
- [11] I.R. Klebanov, I. Kogan and A.M. Polyakov, Phys. Rev. Lett. 71 (1993) 3243
- [12] Reviews of this work can be found in the talks of N. Berkovits and R. Kallosh at Strings '99 (July 1999, Potsdam), available at <http://strings99.aei-potsdam.mpg.de/cgi-bin/viewit.cgi>
- [13] N. Dorey, T. Hollowood, V. Khoze, M. Mattis and S. Vandoren, Nucl. Phys. B552 (1999) 88
- [14] D. Gross, talk at Strings '99 (July 1999, Potsdam), available at <http://strings99.aei-potsdam.mpg.de/cgi-bin/viewit.cgi>
- [15] See *e.g.* A.M. Polyakov, *Gauge fields and strings*, Harwood Academic Publishers (Chur, Switzerland, 1987)
- [16] A. Jevicki and B. Sakita, Nucl. Phys. B185 (1981) 89
- [17] N. Ishibashi and H. Kawai, Phys. Lett. B314 (1993) 190
- [18] G. Horowitz, J. Lykken, R. Rohm and A. Strominger, Phys. Rev. Lett. 57 (1986) 283
- [19] M. Fukuma, N. Ishibashi, H. Kawai and M. Ninomiya, Nucl. Phys. B427 (1994) 139; M. Ikehara, N. Ishibashi, H. Kawai, T. Mogami, R. Nakayama and N. Sasakura, Phys. Rev. D50 (1994) 7467
- [20] N. Ishibashi and H. Kawai, Phys. Lett. B322 (1994) 67; Phys. Lett. B352 (1995) 75
- [21] S. Das and A. Jevicki, Mod. Phys. Lett. A5 (1990) 1639
- [22] G. Moore, N. Seiberg and M. Staudacher, Nucl. Phys. B362 (1991) 665
- [23] A. Jevicki and J. Rodrigues, Nucl. Phys. B421 (278) 1994
- [24] E. Brezin and V. Kazakov, Phys. Lett. B236 (1990) 144; M. Douglas and S. Shenker, Nucl. Phys. B335 (1990) 635; D. Gross and A. Migdal, Phys. Rev. Lett. 64 (1990) 127
- [25] N. Drukker, D. Gross and H. Ooguri, *Wilson loops and minimal surfaces*, hep-th/9904191
- [26] O. Bonnet, J. École Polytech. 19 cah. 32 (1848) 1
- [27] For a recent comprehensive treatment see H. Itoyama and H. Takashino, Prog. Theor. Phys. 97 (1997) 963
- [28] T. Banks, W. Fischler, S. Shenker and L. Susskind, Phys. Rev. D55 (1997) 5112

- [29] M. Srednicki and R. Woodard, Nucl. Phys. B293 (1987) 612
- [30] J.E. Lidsey, *The embedding of superstring backgrounds in Einstein gravity*, hep-th/9907095